

# ***Strobilanthes sunhangii* (Acanthaceae), a new species from Tibet, China**

Jun-Tong Chen<sup>1,2</sup>, Xian-Han Huang<sup>1</sup>, Zhen-Yu Lv<sup>3</sup>,  
Tian-Hui Kuang<sup>1,2</sup>, Jian Luo<sup>4</sup>, Yun-Fei Deng<sup>5</sup>, Tao Deng<sup>1</sup>

**1** Key Laboratory for Plant Diversity and Biogeography of East Asia, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, Yunnan, China **2** University of Chinese Academy of Sciences, Beijing 100049, China **3** School of Life Sciences, Yunnan Normal University, Kunming 650092, Yunnan, China **4** Institute of Plateau Ecology, Tibet Agricultural and Animal Husbandry University, Nyingchi 860000, Tibet, China **5** Key Laboratory of Plant Resources Conservation and Sustainable Utilization, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou 510650, Guangdong, China

Corresponding author: Tao Deng ([dengtao@mail.kib.ac.cn](mailto:dengtao@mail.kib.ac.cn)), Yun-Fei Deng ([yfdeng@scbg.ac.cn](mailto:yfdeng@scbg.ac.cn))

---

Academic editor: E. Fischer | Received 21 September 2020 | Accepted 30 October 2020 | Published 13 November 2020

---

**Citation:** Chen J-T, Huang X-H, Lv Z-Y, Kuang T-H, Luo J, Deng Y-F, Deng T (2020) *Strobilanthes sunhangii* (Acanthaceae), a new species from Tibet, China. PhytoKeys 166: 117–127. <https://doi.org/10.3897/phytokeys.166.58831>

---

## **Abstract**

A new species of Acanthaceae, *Strobilanthes sunhangii*, is described from Mêdog County, Tibet, China. Morphologically, the new species is closely similar to *S. medogensis* and *S. divaricata*, but *S. sunhangii* differs in having glabrous stems, longer spikes, glabrous rachis, double curved corolla and glabrous calyx, different stamens and style.

## **Keywords**

Mêdog, morphological evidence, new species, *Strobilanthes sunhangii*



## Introduction

*Strobilanthes* Blume (1826: 781) is one of the most diverse genera in the family Acanthaceae (Ruellieae: Strobilanthinae) and comprises approximately 450 species mainly distributed in tropical and subtropical regions of Asia with some species extending to Pacific Islands (Hu and Tsui 2002; Hu et al. 2011; Tripp et al. 2013; Chen et al. 2019; Deng 2019). Historically, there has long been disagreement about whether the genus might be adopted on a broad or narrow circumscription (von Esenbeck 1847; Anderson 1867; Clarke 1885; Terao 1983). Beremekap (1944) divided Strobilanthinae into many small genera. However, molecular studies (Moylan et al. 2004) have shown that these segregate genera are nested within *Strobilanthes* and the broad concept of *Strobilanthes* has been generally accepted (Deng et al. 2006; Wood and Scotland 2009; Hu et al. 2011; Wood 2014; Deng 2019). In China, Hu et al. (2011) recorded 128 species mainly distributed in SE to SW China.

China has a vast territory with a wide range of complex and diverse topographies and soils and covering several climate types, which contribute to the wealth of Chinese botanical diversity (Sun et al. 2017; Chen et al. 2018a, b, 2020). Mêdog County, in which the highest peak is Mt. Namjagbarwa (7,782 m) in the north and the lowest point is Pasighat (154 m), has a rich biodiversity, including 1,805 species of seed plants (Sun and Zhou 1996, 2002; Yang and Zhou 2015; Chen et al. 2018b) and some new species have recently been reported from this region, such as *Cornus sunhangii* (Lv et al. 2019).

During a recent expedition in Mêdog, we discovered a species of *Strobilanthes* with very unusual morphological characters. After observations of wild living plants, herbarium specimens, laboratory studies and consultation of relevant literature (Blume 1826; Anderson 1867; Clarke 1885; Li 1985; Wood 1994, 2001, 2014; Carine and Scotland 2002; Hu and Tsui 2002; Deng et al. 2006; Hu et al. 2011; Adhikari 2018; Chen et al. 2019; Thomas et al. 2019), it is concluded that it does not match any other known species of *Strobilanthes* and represents an undescribed species. Therefore, it is described below as *Strobilanthes sunhangii* T. Deng, J.T. Chen & Y.F. Deng.

## Materials and methods

The specimens of *Strobilanthes sunhangii* were collected from Mêdog County in Tibet and studied at the herbarium of Kunming Institute of Botany (KUN). Morphological characters, recorded for the new species, were based on fresh plants and dried specimens. Pollen grains were collected from dry specimens and the operation method for pollen measurement followed Chen et al. (2019). Morphological comparisons of *S. sunhangii*, with the related taxa *S. medogensis* (H.W. Li) J.R.I. Wood & Y.F. Deng and *S. divaricata* (Nees) T. Anderson are provided in Table 1.



## Results

### Taxonomic treatment

***Strobilanthes sunhangii* T. Deng, J.T. Chen & Y.F. Deng, sp. nov.**

urn:lsid:ipni.org:names:77212877-1

Figs 1–4

**Type.** CHINA. Tibet: Mêdog County, Beibeng Town, 29.23319N, 95.17693E, elev. ca. 1,470 m, 5 Oct 2018, *H. Sun, T. Deng & Z.M. Li Sunhang19964* (**holotype**: KUN1345286!; **isotypes**: IBSC!, KUN1345287!, KUN1345288!)

**Diagnosis.** *Strobilanthes sunhangii* resembles *S. medogensis*, but differs by its stems subterete, glabrous (vs. slightly sulcate, bifariously puberulent); spikes (7–)11–22 cm long (vs. 3–6 cm long), rachis glabrous (vs. bifariously pubescent); corolla outside and lobes pinkish-white, inside purplish-pink (vs. corolla yellowish-white, but dull purple on lobes), the tube bent to ca. 90° twice (vs. straight), lobes apices emarginate (vs. rounded); calyx 7–8 mm long, glabrous, 5-lobed to middle (vs. ca. 12 mm long, gland-tipped pubescent, 5-lobed to base), lobes ovate, equal (vs. narrowly oblong-elliptic, subequal); stamens included (vs. exserted), filaments of shorter pair ca. 2 mm long (vs. 7–8 mm long), longer pair ca. 4 mm long (vs. 9–10 mm long), anther thecae ca. 4 mm (vs. ca. 1 mm); style with sparse gland-tipped trichomes (vs. glabrous).


**Description.** Undershrubs 1–2 m high, gregarious. Stems subterete, glabrous, slightly swollen at node. Leaves slightly anisophyllous, smaller one ca. 2/3 of the larger one in size; petiole 3–7 mm long or petiole of smaller leaf in each pair shorter, sulcate, glabrous; blades ovate to lanceolate-ovate, smaller ones 4.2–6.9 × 1.2–2.0 cm, larger ones 8.2–10.1 × 2.2–3.2 cm, both surfaces glabrous, densely covered with numerous linear cystoliths, secondary veins 3–5 on each side of midvein and prominent on both surfaces, base rounded to broadly cuneate, slightly oblique, margin serrulate, apex acuminate and narrowly caudate. Inflorescences of axillary spikes, simple or rarely 2-furcate, slender, (7–) 11–22 cm long; rachis glabrous; bracts and bracteoles not seen, caducous. Flowers (0.6–) 1.0–2.5 (–3.0) cm apart on rachis. Calyx 7–8 mm long, glabrous, 5-lobed to middle; lobes ovate, equal, margins narrowly scarious, 1-nerved abaxially, conspicuous after drying, apices acuminate. Corolla 2.8–3.3 cm long, campanulate, outside and lobes pinkish-white, inside purplish-pink, outside glabrous, inside glabrous except for bifarious hirsute retaining the style (Fig. 3E); tube basally cylindrical, ca. 3 mm wide and ca. 6 mm long, then bent to ca. 90° and gradually widened to 9–12 mm wide at middle and 1.6–1.8 cm long, then secondly bent to ca. 90° and tube upper cylindrical 9–12 mm wide and 1.6–1.8 cm long; lobes broadly elliptic, 8–9 × 7–8 mm, apices emarginate (Fig. 3A, B). Stamens 4, didynamous, included; the united part of filaments glabrous at base, with dense villous on middle and margin, with slightly sparse villous on upper (Fig. 3E), shorter pair ca. 2 mm long, longer pair ca. 4 mm long; anther thecae oblong, ca. 4 mm long. Pollen grains prolate in outline,






**Figure 1.** Photograph of the holotype of *Strobilanthes sunhangii* T. Deng, J.T. Chen & Y.F. Deng (KUN barcode 1345286!).





**Figure 2.** *Strobilanthes sunhangii* T. Deng, J.T. Chen & Y.F. Deng **A** plant showing flowering branch and leaves **B** flower (view from side) **C** pistil, ovary and style **D** calyx **E** opened corolla showing androecium (Drawn by Xiaoshuang Zhang based on the holotype of Sunhang19964).





**Figure 3.** *Strobilanthes sunhangii* T. Deng, J.T. Chen & Y.F. Deng **A** habitat and flowering branch **B, C** flowering branch and flower (view from front) **D, E** flower (view from side) **F** calyx **G** style **H** leaf adaxial surfaces **I** leaf abaxial surfaces.


74.1 (72.2–76.1)  $\times$  49.7 (48.1–50.4)  $\mu\text{m}$ , P/E = 1.49, tricolporate, ribbed pseudocolpi 12, exine reticulate (Fig. 4). Ovary ca. 5 mm long, glabrous; style ca. 1.8 cm long, with sparse gland-tipped trichomes. Capsule not seen.

**Phenology.** Flowering from June–October.

**Etymology.** *Strobilanthes sunhangii* is named after Prof. Hang Sun (1963–) for his outstanding contributions to the flora of Himalayan regions and collecting this new species for the first time. Verucular name: The Chinese name is given as “双曲马蓝” (shuāng qū mǎ lán), referring to the double-curved corolla of the new species.

**Distribution and habitat.** The new species is currently known only from Mêdog County, Tibet, China. It grows in the moist evergreen forest at an elevation of 1,200–1,800 m. The associations include *Exbucklandia populnea* (R. Br.) R. W. Brown (Hamamelidaceae), *Viburnum* sp. (Viburnaceae), *Rubus* sp. (Rosaceae), *Campanumoea inflata* (Hook. f. & Thomson) C.B. Clarke (Campanulaceae), *Tripterospermum volubile* (D. Don) H. Hara (Gentianaceae) and *Impatiens* sp. (Balsaminaceae), amongst others.





**Figure 4.** Pollen grains of *Strobilanthes sunhangii* T. Deng, J.T. Chen & Y.F. Deng **A** equatorial view **B** polar view **C** ornamentation.

**Conservation status.** The new species was only discovered in Beibeng Town, Mêdog County. About 1,000 individuals were observed and the extent of occurrence was ca. 5 km<sup>2</sup>. Further explorations to obtain the precise population status are needed to assess its conservation status. Based on available data, the new species is assigned to the category ‘Data Deficient’ (DD) of the International Union for Conservation of Nature (IUCN 2019).

**Specimens of *S. medogensis* examined.** – CHINA. **Tibet:** Mêdog County, Gedang Town, elev. ca. 2,400 m, 27 Aug 1974, *QTP Expedition 74-4911* (holotype: KUN barcode 1219209!); Mêdog County, Gedang Town, 2,450 m elev., 11 Sept 1982, *B.S. Li & S.Z. Cheng 814* (PE); Mêdog County, Gedang Town, 1,940 m elev., 13 Oct 1982, *S.Z. Cheng & B.S. Li 1457* (PE); Mêdog County, 1,100 m elev., 1 Dec 1982, *S.Z. Cheng & B.S. Li 3238* (PE); Nyingchi County, Pailong Town, 2,175 m elev., 8 Sept 2006, *Tiber-MacArthur 261* (A); Lung County, Chayne chu, 2,750 m elev., 10 Jul 1936, *Ludlow & Sherriff 2336* (BM, E).

**Specimens of *S. divaricata* examined.** – INDIA. **Meghalaya:** Khasia, 1,524 m elev., 5 Oct 1886, *C. B. Clarke 45219* (BM, CAL, LE); Khasia, 1,219 m elev., 12 Nov 1871, *C. B. Clarke 15479* (CAL, K, LE); Khasia, Oct 1850, *Simons 482* (CAL). **Sikkim:** Khuping, 1,676 m elev., 26 Sept 1884, *C. B. Clarke 35904* (LE); Khuping, 1,676 m elev., 26 Sept 1884, *C. B. Clarke 35920* (LE). – NEPAL. Koshi Zone, Sankhuwasabha District, 2,020 m elev., 16 Aug 1998, *S. Noshiro et al. 9840046* (A); Dhawalagiri Zone, Myagdi District, 1,820–2,360 m elev., 10 Sept 1996, *M. Mikage et al. 9681329* (A).

## Discussion

Morphologically, the new species is closely related to *Strobilanthes medogensis* in having similar leaves and slender spikes, but differs in having glabrous stems, longer spikes, glabrous rachis, double curved corolla and glabrous calyx, different stamens and style (Li 1985; Wood 1994; Hu and Tsui 2002; Deng et al. 2006; Hu et al. 2011). In addition, the new species is also related to *S. divaricata* in the glabrous stems and calyx, slender spikes, but differs in having longer spikes, double curved corolla, different stamens and style (Wood 2001; Adhikari 2018). The new species is quite different from all other known species in its unique corolla shape. The corolla of *S. sunhangii* are bent twice, at the base and middle of throat, respectively, while that of other species is



**Table 1.** Morphological comparisons of *Strobilanthes sunhangii* with related species.

Characters	<i>S. sunhangii</i>	<i>S. medogensis</i>	<i>S. divaricata</i>
Stems	Stems subterete, glabrous.	Stems slightly sulcate, bifariously puberulent.	Stems often zigzag above, angular, glabrous.
Leaves	Leaves slightly anisophyllous, leaf blade ovate to lance-ovate, smaller of pair ca. 2/3 size of larger one, 4.2–6.9 × 1.2–2.0 cm and larger ones 8.2–10.1 × 2.2–3.2 cm, base rounded to broadly cuneate, both surfaces glabrous and with numerous linear cystoliths.	Leaves anisophyllous, leaf blade ovate, smaller of pair ca. 2/3 size of larger one, (1.9–)4.0–6.0 × (0.5–)1.8–2.1 cm and larger ones 5.8–9.0 × 2.1–3.7 cm, base rounded to subcordate, both surfaces glabrous, adaxially with numerous cystoliths.	Leaves strongly anisophyllous, smaller leaves ca. 1/3 length of larger leaves, ovate, 1.5–6 × 0.5–2 cm, base rounded, larger ones lanceolate to elliptic, 8–18 × 2–5 cm, base cuneate or attenuate to a short petiole, both surfaces glabrous.
Inflorescences	Spikes slender, (7–)11–22 cm; rachis glabrous.	Spikes 3–6 cm; rachis bifariously pubescent.	Spikes 1–6 cm, rachis glabrous to sparsely pilose.
Corolla colour	Corolla outside and lobes pinkish-white, inside purplish-pink.	Corolla yellowish-white but dull purple on lobes.	Corolla purple.
Corolla	Corolla 2.8–3.3 cm long, tube basally cylindrical and ca. 3 mm wide by ca. 6 mm long, then bent to ca. 90° and gradually widened to 9–12 mm wide at middle by 16–18 mm long, then second bent to ca. 90° and tube upper cylindrical and 9–12 mm wide by 16–18 mm long.	Corolla ca. 2.5 cm long, straight, tube basally cylindrical and ca. 2 mm wide by ca. 1 cm long, then gradually widened to 1–1.5 cm at mouth.	Corolla 3–3.5 cm long, tube cylindrical at base for 1 cm, then gradually widened up to 0.8 cm at mouth, straight or slightly bent towards mouth.
Corolla lobes	Corolla lobes widely elliptic, ca. 8–9 × 7–8 mm, apices emarginate.	Corolla lobes broadly ovate, ca. 3 × 8 mm, apices rounded.	Corolla lobes 5–5.5 × 6 mm, apices obtuse or rounded.
Calyx	Calyx 7–8 mm, glabrous, 5-lobed to middle; lobes ovate, equal, margins narrowly scarious.	Calyx ca. 12 mm, gland-tipped pubescent, 5-lobed to base; lobes narrowly oblong-elliptic, subequal.	Calyx 8–15 mm, glabrous rarely pilose at tips, lobes linear-lanceolate, acuminate, subequal.
Stamens	Stamens included; filaments of shorter pair ca. 2 mm, longer pair ca. 4 mm; anther thecae oblong, ca. 4 mm.	Stamens exerted; filaments of shorter pair 7–8 mm, longer pair 9–10 mm; anther thecae ca. 1 mm, incurved.	Stamens included, filaments of shorter pair incurved, 1 mm, longer pair unequal, longer one 5–5.5 mm, shorter one 4–4.5 mm; anther thecae ca. 1 mm.
Style	Style ca. 1.8 cm, with sparse gland-tipped trichomes.	Style 2.7–2.8 cm, glabrous.	Style 2–3 cm, sparsely glandular trichomes at base.
Distribution	Tibet (Mêdog), China.	Tibet (Mêdog), China.	India (NE-India, Darjeeling, Sikkim); Bhutan; Nepal.

straight or curved at the middle of the throat. The detailed morphological comparisons amongst *S. sunhangii*, *S. medogensis* and *S. divaricata* are shown in Table 1.

Till now, no infrageneric classification has been established. Bremekamp (1944) divided the subtribe Strobilanthinae into 54 genera and 27 informal groups. However, his classification is not supported by the studies of the only molecular phylogeny of *Strobilanthes* (Moylan et al. 2004) and morphological analysis (Carine and Scotland 2002). In this study, *S. divaricata* and *S. medogensis*, two species closest to the new species, were not included. Bremekamp placed *S. divaricata* into his new genus *Diffugossa* Bremek., which is characterised by the single flowers on each node on rachis, arranged in an open panicle, bracts small, deciduous and corolla straight. However, *S. divaricata* is quite different from other species of *Diffugossa*, which was informally placed in Panicle groups by Wood and Scotland (2003), in its paired flower on the rachis, arranged in spikes. *S. medogensis* was placed in *Goldfussia* Nees, which is characterised by the flowers arranged in congested head-like spikes and then compounded into small panicles, stamens 2-paired, the shorter pair unequal, nodding, anthers spherical (Li 1985). Obviously,



these three species form a species complex and further studies on the placement in the genus are necessary, depending on the establishment the infrageneric classification.

Pollen morphology has been widely used in species delimitation of *Strobilanthes* (Chen et al. 2019), so we observed and measured the pollen grains of this new species. Pollens grains of the new species are prolate,  $74.1 (72.2\text{--}76.1) \times 49.7 (48.1\text{--}50.4) \mu\text{m}$ ,  $P/E = 1.49$ , 3-colporate, 12-pseudocolpate, with reticulate exine. The detailed pollen grains morphology is provided in Fig. 4. The pollen grains of *S. medogensis* (as *S. campaniformis*) were observed by Wood (1994) and those of *S. divaricata* have not yet been observed. Both *S. sunhangii* and *S. medogensis* belong to type 3 recognised by Hu et al. (2011). Therefore, the close relationship between *S. sunhangii* and *S. medogensis* is also supported by the pollen morphology.

This species was observed by the authors during the Mêdog expedition in 2018. To date, we only collected flowering specimens and whether this new species has a plietesial life history requires further explorations.

## Acknowledgements

We are grateful to Ms. Xiao-shuang Zhang for her excellent illustration of *Strobilanthes sunhangii*. This study was supported by grants from the Second Tibetan Plateau Scientific Expedition and Research (STEP) program (2019QZKK0502), the National Natural Science Foundation of China-Yunnan joint fund to support key projects (U1802232), the Major Program of the National Natural Science Foundation of China (31590823), the National Natural Science Foundation of China (31700165, 31270247), the Strategic Priority Research Program of Chinese Academy of Sciences (XDA20050203), the National Key R & D Program of China (2017YFC0505200), the Youth Innovation Promotion Association of the Chinese Academy of Sciences (2019382), the Young Academic and Technical Leader Raising Foundation of Yunnan Province (2019HB039) and the Chinese Academy of Sciences “Light of West China” Program.

## References

- Adhikari B (2018) *Strobilanthes*. In: Watson MF, Akiyama S, Ikeda H, Pendry C, Rajbhandari KR, Shrestha KK (Eds) *Flora of Nepal Acanthaceae*: Web edition 2. Royal Botanic Garden Edinburgh, Edinburgh, 36–49. <http://www.floraofnepal.org/accountlibrary?name=Acanthaceae>
- Anderson T (1867) An Enumeration of the Indian species of Acanthaceae. *Journal of the Linnean Society of London. Botany* 9: 425–526. <https://doi.org/10.1111/j.1095-8339.1867.tb01309.x>
- Bremekamp CEB (1944) Materials for a Monograph of the Strobilanthinae. *Verhandelingen der Nederlandsche Akademie van Wetenschappen, Afdeling Natuurkunde* 2 41(1):1–306.



- Blume CL (1826) Bijdragen tot de Flora van Nederlandsch Indië 14. Ter Lands Drukkerij, Batavia, 731–850. <https://doi.org/10.5962/bhl.title.395>
- Carine MA, Scotland RW (2002) Classification of Strobilanthes (Acanthaceae): Trying to classify the unclassifiable? *Taxon* 51(2): 259–279. <https://doi.org/10.2307/1554897>
- Chen JT, Zhong J, Shi XJ, Zhang QX, Sun M (2018a) *Chrysanthemum yantaiense*, a rare new species of the Asteraceae from China. *Phytotaxa* 374(1): 92–96. <https://doi.org/10.11646/phytotaxa.374.1.9>
- Chen YS, Deng T, Zhou Z, Sun H (2018b) Is the east Asian flora ancient or not? *National Science Review* 5(6): 920–932. <https://doi.org/10.1093/nsr/nwx156>
- Chen FL, Deng YF, Xiong ZB, Ran JC (2019) *Strobilanthes hongii*, a new species of Acanthaceae from Guizhou, China. *Phytotaxa* 388(1): 135–144. <https://doi.org/10.11646/phytotaxa.388.1.7>
- Chen JT, Zhang DG, Lv ZY, Huang XH, Liu PJ, Yang JN, Yang JY, Tojibaev K, Deng T, Sun H (2020) *Oxytropis shennongjiaensis* (Fabaceae), a new species from Hubei, Central China. *PhytoKeys* 149: 117–128. <https://doi.org/10.3897/phytokeys.149.49533>
- Clarke CB (1885) Acanthaceae. In: Hooker JD (Ed.) *Flora of British India* 4. Reeve & Co., London, 387–558.
- Deng YF (2019) Transfer of the Philippine species of *Hemigraphis* Nees to *Strobilanthes* Blume (Acanthaceae). *Phytotaxa* 404(5): 1–3. <https://doi.org/10.11646/phytotaxa.404.5.3>
- Deng YF, Wood JRI, Scotland RW (2006) New and reassessed species of *Strobilanthes* (Acanthaceae) in the flora of China. *Botanical Journal of the Linnean Society* 150(3): 369–390. <https://doi.org/10.1111/j.1095-8339.2006.00473.x>
- Hu CC, Tsui HP (2002) Acanthaceae. In: Hu CC (Ed.) *Flora Reipublicae Popularis Sinicae* (Vol. 70). Science Press, Beijing, 309 pp. [in Chinese]
- Hu JQ, Deng YF, Wood JRI (2011) *Strobilanthes*. In: Wu ZY, Raven PH, Hong DY (Eds) *Flora of China* vol. 19. Beijing & Missouri Botanical Garden Press, St. Louis, 381–429.
- IUCN (2019) Guidelines for Using the IUCN Red List Categories and Criteria. version 13. Prepared by the Standards and Petitions Subcommittee of the IUCN Species Survival Commission, 113 pp. <http://cmsdocs.s3.amazonaws.com/RedListGuidelines.pdf>
- Li HW (1985) Acanthaceae. In: Wu ZY (Ed.) *Flora Xizangica* (Vol. 4). Science Press, Beijing, 411–416. [in Chinese]
- Lv ZY, Huang XH, Zhang X, Yusupov Z, Wang HC, Tojibaev K, Deng T, Li ZM (2019) *Cornus sunhangii* (Cornaceae), a new species from Tibet (China). *Phytotaxa* 409(5): 273–282. <https://doi.org/10.11646/phytotaxa.409.5.3>
- Moylan EC, Bennett JR, Carine MA, Olmstead RG, Scotland RW (2004) Phylogenetic relationships among *Strobilanthes* s.l. (Acanthaceae): Evidence from ITS, nrDNA, TrnL-F cpDNA, and morphology. *American Journal of Botany* 91(5): 724–735. <https://doi.org/10.3732/ajb.91.5.724>
- Sun H, Zhou ZK (1996) The characters and origin of the flora from the Big Bend Gorge of Yalu Tsangpo (Brahmabutra) River, Eastern Himalayas. *Yunnan Zhi Wu Yan Jiu* 15: 185–204. [in Chinese]
- Sun H, Zhou ZK (2002) Seed plants of the Big Bend Gorge of Yalu Tsangpo (Brahmabutra) River in SE Tibet, E Himalayas. Yunnan Science and Technology Press, Kunming, 3–46. [in Chinese]



- Sun H, Zhang JW, Deng T, Boufford DE (2017) Origins and evolution of plant diversity in the Hengduan Mountains, China. *Plant Diversity* 39(4): 161–166. <https://doi.org/10.1016/j.pld.2017.09.004>
- Terao H (1983) Taxonomic study of the genus *Strobilanthes* Blume (Acanthaceae) generic delimitation and infrageneric classification. PhD thesis. Kyoto University, Japan.
- Thomas S, Mani B, Britto SJ, Pillai PAVK (2019) *Strobilanthes tricostata*, a new species of Acanthaceae from the Western Ghats, India. *Phytotaxa* 413(3): 244–250. <https://doi.org/10.11646/phytotaxa.413.3.5>
- Tripp EA, Daniel TF, Fatimah S, McDade LA (2013) Phylogenetic relationships within Ruelieae (Acanthaceae) and a revised classification. *International Journal of Plant Sciences* 174(1): 97–137. <https://doi.org/10.1086/668248>
- von Esenbeck N (1847) Acanthaceae. In: de Candolle AP (Ed.) *Prodromus systematis naturalis regni vegetabilis* 11. Masson, Paris, 46–519.
- Wood JRI (1994) Notes relating to the flora of Bhutan: XXIX. Acanthaceae, with special reference to *Strobilanthes*. *Edinburgh Journal of Botany* 51(2): 175–274. <https://doi.org/10.1017/S0960428600000871>
- Wood JRI (2001) Acanthaceae. In: Grierson AJC, Long DG (Eds) *Flora of Bhutan: including a record of plants from Sikkim* 2 (3). Royal Botanical Garden, Edinburgh & Royal Government of Bhutan, 1243–1293.
- Wood JRI (2014) New names and combinations in Indian Acanthaceae. *Novon* 23(3): 385–395. <https://doi.org/10.3417/2013046>
- Wood JRI, Scotland RW (2003) *Strobilanthes*: Panicked Species from East Asia. *Kew Bulletin* 58(3): 679–702. <https://doi.org/10.2307/4111148>
- Wood JRI, Scotland RW (2009) New and little-known species of *Strobilanthes* (Acanthaceae) from India and South East Asia. *Kew Bulletin* 64(1): 3–47. <https://doi.org/10.1007/s12225-009-9098-2>
- Yang N, Zhou XW (2015) *Plants of Mêdog*. China Forestry Press, Beijing, 34–79. [in Chinese]